

XMM-NEWTON OBSERVATIONS OF THE EXTREMELY LOW ACCRETION RATE POLARS SDSSJ155331.12+551614.5 AND SDSSJ132411.57+032050.5

PAULA SZKODY¹, LEE HOMER¹, BING CHEN², ARNE HENDEN³, GARY D. SCHMIDT⁴, SCOTT F. ANDERSON¹, D. W. HOARD⁵,
WOLFGANG VOGES⁶, J. BRINKMANN⁷

Draft version February 2, 2008

ABSTRACT

XMM-Newton observations of the Polar SDSSJ155331.12+551614.5 reveal that all the X-ray flux emerges at energies < 2 keV. The best fit to the spectrum is with a thermal plasma with $kT = 0.8$ keV plus a 20-90 eV black body, yielding a thermal X-ray luminosity of $8.0\text{--}9.5 \times 10^{28}$ ergs s⁻¹. The low temperature and X-ray luminosity, together with the lack of variation of the X-ray flux during the observations, are all consistent with an extremely low accretion rate that puts the system in the bombardment regime of accretion, rather than accretion involving a standoff shock. It is likely that the observed X-rays originate from the M dwarf secondary star, thus providing a base activity level for late main sequence stars in close binaries. SDSSJ132411.57+032050.5 is detected by *XMM-Newton* at the faint EPIC-pn count rate of 0.0012 ± 0.0003 giving an upper limit to the X-ray luminosity of $\sim 7 \times 10^{28}$ ergs s⁻¹ for a distance of 300 pc, which is also consistent with the above scenario.

Subject headings: cataclysmic variables — stars:individual(SDSSJ155331.12+551614.5, SDSSJ132411.57+032050.5) — stars:magnetic fields — X-rays:stars

1. INTRODUCTION

The early results from the Sloan Digital Sky Survey (SDSS; York et al. 2000, Abazajian et al. 2003, 2004) have shown that this survey is very effective in finding the shortest period (oldest) and lowest mass transfer rate (faintest) cataclysmic variables (Szkody et al. 2002; 2003a). Included among the new discoveries are two systems containing magnetic white dwarfs (i.e. Polars) with extreme cyclotron features; that is, very narrow, highly polarized cyclotron emission harmonics that show dramatic changes during the orbital cycle (Szkody et al 2003b). The lack of strong Balmer emission lines and the narrow appearance of the cyclotron features imply very low temperature plasmas (< 1 keV) and accretion rates of $< 10^{-13} M_{\odot}$ yr⁻¹ (Ferrario et al. 2002). In this regime, the energy of the incoming flow is dissipated through particle collisions in the white dwarf atmosphere rather than in an accretion shock (Wickramasinghe & Ferrario 2000), a situation which is sometimes termed a “bombardment solution” (Kuijpers & Pringle 1982).

In order to check the consistency of this model, we obtained X-ray measurements of these two systems SDSSJ155331.12+551614.5 and SDSSJ132411.57+032050.5 (hereafter abbreviated as SDSSJ1553 and SDSSJ1324 for simplicity) with the *XMM-Newton* satellite. From the data shown in Szkody et al. (2003b), these objects are known to have orbital

periods of 4.39 hr and 2.6 hr respectively, derived from the > 1 mag sinusoidal modulations in the V and R light curves created by the changing cyclotron harmonics. The sinusoidal shapes of the variations imply that the magnetic poles are always in view. In each system, the wavelengths of the observed cyclotron harmonics are consistent with the 3rd and 4th harmonics in a magnetic field near 60 MG. The presence and strength of TiO bands in SDSSJ1553 imply a secondary star of spectral type near M5V, and assuming a main-sequence mass-radius relation, a corresponding distance of 100 pc. The same assumptions for an M6V star in SDSSJ1324 yield a distance of $\sim 300\text{--}500$ pc.

Detailed modelling of the spectrum of SDSSJ1553 (Ferrario et al. 2002) indicates that it has an accretion rate three orders of magnitude smaller than a typical Polar system, and an underlying very cool white dwarf with temperature of only 5000-8000K. With these very unusually low accretion rates and cool white dwarfs, X-rays are not expected from the accretion spots (Woelk & Beuermann 1992). The ROSAT All Sky Survey provides upper limits of 0.04 and 0.02 c s⁻¹ for SDSSJ1553 and SDSSJ1324.

A recent survey of 37 Polars using *XMM-Newton* showed that 16 were in a low or reduced accretion state (Ramsay et al. 2004), with 6 not detected. Of the 10 with detections, 8 showed orbital variations, indicating that some accretion was still occurring. Of the 2 systems with no orbital variation (CP Tuc and MN Hya), the observation times were too short to provide a spectrum. However, by assuming a 1 keV thermal plasma and an absorption of 1×10^{20} cm⁻², Ramsay et al. (2004) inferred X-ray luminosities for the two systems that were 15-20 times greater than that expected for normal late main sequence stars. With our long observation times on the extremely low accretion systems SDSSJ1553 and SDSSJ1324, we are able to accomplish a spectral fit for SDSSJ1553 and obtain better luminosity constraints for the low states in both Polars.

¹ Department of Astronomy, Box 351580, University of Washington, Seattle, WA 98195

² XMM-Newton Science Operations Centre, ESA/Vilspa, 28080, Madrid, Spain

³ Universities Space Research Association/US Naval Observatory, Flagstaff Station, P. O. Box 1149, Flagstaff, AZ 86002-1149

⁴ The University of Arizona, Steward Observatory, Tucson, AZ 85721

⁵ Spitzer Science Center, California Institute of Technology, 1200 East California Boulevard, Pasadena, CA 91125

⁶ Max-Planck-Institut für Extraterrestrische Physik, Geisenbachstr. 1, D-85741 Garching, Germany

⁷ Apache Point Observatory, P. O. Box 59, Sunspot, NM 88349-0059

2. OBSERVATIONS

The *XMM-Newton* satellite observes with three X-ray cameras (the EPIC-pn has about twice the sensitivity of either the EPIC-MOS1 or MOS2; Turner et al. 2001) as well as X-ray reflection grating spectrographs (our two systems were too faint for any useful data from the spectrographs) and an optical monitor (OM; Mason et al. 2001). The OM was used with a V filter for SDSSJ1553 and a white light filter for SDSSJ1324. The data were analyzed following the ABC guide from the *XMM-Newton* US GOF⁸ and analysis threads from the main Vilspa⁹ websites, using the SAS v6.0 and calibration files current to 2004 March 23. This included reprocessing of ODF files to produce new event files. These were then screened using the standard expressions, and energies were restricted to the range 0.1-10 keV. Finally, revised good time intervals were derived from high energy background light curves to exclude intervals of high background flaring. For the pn, source data were extracted using a 320 pixel radius circular aperture (this only encircles $\sim 70\%$ of the energy, but reduces the background contribution for this low count-rate source) and background was selected from adjacent rectangular regions at comparable detector Y locations as the target. As advised in the guide, a conservative choice of event selection, pattern = 0, was adopted. For the two MOS detectors, a similar aperture size was used, but the background was determined from an annulus surrounding the source, chosen to exclude a neighboring faint source, and an event pattern ≤ 12 was also applied.

Light curves for both source and background were extracted using the SAS task `evselect`, employing the same extraction regions as described above, but with a less conservative pattern ≤ 4 for the pn events. The task `lccor` scaled and applied background subtraction, dead-time and vignetting corrections, to produce net source light curves with 100s binning. Finally, the time stamps were converted from JD(TT) to HJD(UT), using `FTOOLS`¹⁰ tasks.

The X-ray observation of SDSSJ1553 began on 2003 February 17 but was halted prematurely due to high radiation levels. Additional data were obtained on 2003 March 12 to complete the observation. The OM provided useful data during the X-ray observations (Figure 1), while higher time resolution, ground-based CCD differential photometry in V and open filters was obtained on 2003 February 19 and 22 and June 29 using the United States Naval Observatory (USNO) 1m telescope (Figure 2). The light curves all show the same variations as present in Szkody et al (2003b), indicating that the accretion state was similarly low. The variation in the V and open filters is caused by the changing strength of the $n=3$ cyclotron harmonic at 6200Å. In addition, the ground-based and OM optical light curves allow a relative phasing of the X-ray data. Due to the high radiation levels and other screening, the good time intervals of the X-ray data only span portions of the total observation length. With phase 0 defined as the minimum of the optical variation, the February good time intervals cover phases 0.75-1.43 (87.61-87.73 in Figure 1) and




FIG. 1.— OM V light curves of SDSS J1553 during the February and March *XMM-Newton* observations. Data are binned into 10 minute segments with individual 1σ errors of 0.06 mag.

the March observation covers phases 0.88-1.49 (111.24-111.35 in Figure 1). Thus, most of the orbit is covered, and the portions from minimum to maximum light are sampled in each X-ray observation.

SDSSJ1324 was observed with *XMM-Newton* on 2004 Jan 25. The source was barely detected with the most sensitive EPIC-pn instrument, so no time resolution was possible. The OM data with 10 min bins and the USNO data from the night before the *XMM-Newton* observations (Figure 3) show the 2.6 hr variation previously identified as the orbital period. The X-ray observations of both systems are summarized in Table 1.

3. RESULTS

3.1. SDSSJ1553

The count rates for the February and March datasets on SDSSJ1553 are consistent with each other, and the binned X-ray light curves show no variation throughout the observation intervals. A plot of X-ray count rate vs energy (Figure 4) for the EPIC-pn, -MOS1 and -MOS2 data shows that all of the observed X-ray flux for SDSSJ1553 emerges below 2 keV. A joint fit was performed in XSPEC¹¹ on all the pn and MOS data from each of the two runs. We binned the data to give > 5 counts per bin and used Cash statistics to find the best overall fit. To accommodate any discrepancies in the relative pn and MOS effective area calibrations, the normalizations were allowed to vary. Similarly, the normal-

⁸ <http://heasarc.gsfc.nasa.gov/docs/xmm/abc/abc.html>

⁹ <http://xmm.vilspa.esa.es/external/xmm-sw-cal/sas.shtml>

¹⁰ <http://heasarc.gsfc.nasa.gov/lheasoft/ftools/>

¹¹ <http://heasarc.gsfc.nasa.gov/docs/xanadu/xspec/index.html>

izations between the February and March runs were allowed to vary. SDSSJ1553 is nearby ($d = 100$ pc) and at a Galactic latitude of 47 deg, so the column density to the source is expected to be low. Hence, this parameter was fixed at $1.3 \times 10^{20} \text{ cm}^{-2}$, the value obtained from the HEASARC column density tool¹².

Because of the uncertainties in the calibration at the lowest energies, fits were attempted using energies >0.15 keV for pn and >0.2 keV for MOS, as well as more conservative limits of >0.3 keV (pn) and >0.5 keV (MOS). A thermal plasma (mekal) model with $kT=0.78 \pm 0.03$ keV provided the best fit in both cases. However, Figure 4 shows that there may be an additional soft component present at energies <0.3 keV. This can be fit with a black body of roughly 46 eV which is poorly constrained (20–90 eV is possible). Table 2 gives the parameters of this type of fit and it is shown in Figure 4. We note that there are larger residuals to the fit at around 0.8 keV, most likely due to a combination of the noise and low energy resolution of the data and the complexity of the (unresolved) emission lines in this region (see Mukai et al. (2003) for higher resolution X-ray spectra of cataclysmic variables). The unabsorbed total flux (0.2–10 keV) for the mekal fit alone is $6.7\text{--}7.9 \times 10^{-14} \text{ ergs cm}^{-2} \text{ s}^{-1}$. Using the distance determined from the observed M star (100 pc) yields an X-ray luminosity of $8.0\text{--}9.5 \times 10^{28} \text{ ergs s}^{-1}$.

The lack of hard X-rays and absence of modulation in the light curve corroborate that the X-ray source is not associated with an accretion shock at the magnetic pole, and that this low accretion rate is within the bombardment regime. However, the origin of X-ray flux that emerges even at these low rates is unclear. Ramsay et al. (2004) concluded that there was still some level of ongoing accretion even in the systems showing no orbital modulation. However, this conclusion was based on assumptions about temperature and absorption, and a lack of information about the exact accretion level of CP Tuc and MN Hya. The pn count rates for CP Tuc and MN Hya are comparable to SDSSJ1553, even though the former are 2–5 times further away. This alone implies that they have a higher level of accretion than SDSSJ1553.

Many Polars display a variety of low states, corresponding to different levels of accretion. The prototype, AM Her, is one such example, exhibiting high states with V mag of 12.5, intermediate states near 13.5 and low states near 15.2. Fabbiano (1982) observed AM Her at an intermediate state of 14.5 and found the usual high state components (13.5 keV from the shock and 30 eV from the heated white dwarf) replaced by a single 9 keV component. De Martino et al. (1998) observed AM Her in one of its lowest states ($V=15.1$) and found thermal X-ray flux with a temperature near 5.8 keV for part of the observation and then a drop in X-ray flux by a factor of 7 to the lowest level yet seen. This lowest portion showed a temperature >3.6 keV, with a corresponding luminosity of $3 \times 10^{30} \text{ ergs s}^{-1}$ and an accretion rate of $\geq 4 \times 10^{-13} M_{\odot}/\text{yr}$. Since this temperature is higher than expected for coronal emission (which is usually in the 0.25–1.7 keV range), they concluded that there was likely still some residual accretion, but that the secondary could have a substantial contribution to

the X-ray flux.

SDSSJ1553 has the lowest accretion level known (3 orders of magnitude below most Polars and an order of magnitude below AM Her in its lowest state), a value that is consistent with wind accretion from the companion. There is no broad line emission, nor evidence of any type, for a mass transfer stream from the secondary to the white dwarf. Thus, this system might best serve as a template for the base level of X-ray flux in a Polar.

The extremely low white dwarf temperature of SDSSJ1553 indicates that the average accretion rate over about the last million years (the Kelvin-Helmholtz time scale of the compressionally heated envelope) cannot have been larger than $\sim 3 \times 10^{-12} M_{\odot} \text{ yr}^{-1}$ (Townsend & Bildsten 2004). This raises the distinct possibility that the secondary in SDSSJ1553 has not yet contacted its Roche lobe, a fact consistent with the inferred M5 spectral type if the main-sequence mass-radius relation applies. As such, SDSSJ1553 would be the first pre-Polar known, and possibly a precursor to objects such as RXJ1313.2-3259 (Gänsicke et al. 2000).

The temperature of the plasma that is emitting X-rays in SDSSJ1553 is in the regime for a coronal source. For its orbital period and spectral type, the mass of the secondary is likely near $0.4 M_{\odot}$ (Warner 1995). From the mass-luminosity relation of Malkov, Piskunov & Shipil’kina (1997), the total luminosity of the secondary should be near $1 \times 10^{32} \text{ ergs s}^{-1}$. Given that the secondary should be in the saturated regime with $L_x/L_{bol} < 10^{-3}$, the X-ray luminosity should be less than $10^{29} \text{ ergs s}^{-1}$. The observed X-ray luminosity is compatible with this limit. Thus, SDSSJ1553 appears to be an excellent candidate for coronal emission of X-rays.

Even though the infalling gas cools almost exclusively through electron cyclotron emission at these very low accretion rates, a blackbody component is expected due to surface heating from the downward-directed cyclotron flux. The “bombardment” calculations of Woelk & Beuermann (1992, 1993) indicate that irradiation occurs from a Rosseland optical depth $\tau \approx 10^{-3}$ to 10^{-4} and that reprocessing from the underlying photosphere should appear as an EUV component with a temperature of $\sim 10^5$ K. The spectral fit with a mekal+blackbody model gives a total 0.01–10 keV flux of $1.2\text{--}1.5 \times 10^{-13} \text{ ergs cm}^{-2} \text{ s}^{-1}$, with the blackbody contributing 32–35% of this flux. This is $\sim 25\%$ of the observed cyclotron flux (Ferrario et al. 2002), and for a distance of 100 pc amounts to a luminosity of $4.7\text{--}6.2 \times 10^{28} \text{ ergs s}^{-1}$. Our temperature estimate of 50 eV (600,000 K) leads to an unrealistically small radius for the emission region of $5 \times 10^4 \text{ cm}$, or 0.006% of the radius of a $0.7 M_{\odot}$ white dwarf. Unfortunately, the upturn in the flux occurs at energies where the calibrations are most uncertain, so the temperature is very uncertain and the mere existence of this component must be confirmed (e.g. with more sensitive/better calibrated X-ray observations). Ramsay et al. (2004) used the UV filters on their OM observations to determine that a low temperature component was present outside the bandpass of the XMM detectors. But since their objects exhibited higher accretion rates and X-ray luminosities, it is difficult to directly compare results.

¹² <http://heasarc.gsfc.nasa.gov/cgi-bin/Tools/w3nh/w3nh.pl>




FIG. 2.— Ground-based optical light curves of SDSS J1553 on February 19 and 22 and June 29, which show the same orbital modulations as apparent in Szkody et al. (2003a). February 19 data have 1σ error bars of 0.08 mag, February 22 of 0.05 mag, and June 29 of 0.01 mag.

SDSSJ1324 was too faint to provide more than an upper limit to the X-ray flux. In the pn, the background contribution and faint source counts are comparable within our analysis aperture. A very crude “spectrum” was constructed from 4 energy bins, and a back-

ground power law model was scaled to contribute 50% in the source aperture and then added to the source model. The column found from the HEASARC tool ($2 \times 10^{20} \text{ cm}^{-2}$) was then fixed and the temperature was stepped from 0.2-1 keV. The best fit was at 0.3 keV, but for an

TABLE 1
SUMMARY OF *XMM-Newton* OBSERVATIONS

SDSSJ	Date	UT	PN(c/s)	PN(s)	MOS1(c/s)	MOS1(s)	MOS2(c/s)	MOS2(s)	OM(s)
1553	2003-02-17	02:29-07:38	0.020±0.002	5130	0.009±0.001	6396s	0.006±0.001	6394s	16700
1553	2003-03-12	17:39-20:23	0.026±0.003	7018	0.010±0.001	9085	0.007±0.001	9192	6000
1324	2004-01-25	02:27-07:28	0.0012±0.0003	16234					13824




FIG. 3.— OM white light curve of SDSS J1324 binned by 10 minutes during the January *XMM-Newton* observation (top) and white light curve from USNO the night before the X-ray observation (bottom). USNO data points have 1σ errors ranging from 0.07 at 20.5 mag to 0.15 at 21.5 mag (except for data at times 28.96–28.97, which have errors up to 1 mag because of passing clouds).

TABLE 2
SPECTRAL FITS

SDSSJ	Model	kT_{mek} (keV)	N_H^a	kT_{bb} (eV)	F_{unabs}^a
1553	Mekal	0.78±0.026	1.3		6.7-7.9
1553	Mekal+BB	0.79±0.026	1.3	46	12-15
1324	Mekal	0.2	2.0		<0.6

^aUnits for N_H are 10^{20} cm^{-2} . F is unabsorbed flux over energies of 0.2–10 keV for mekal and 0.01–10 keV for mekal+BB. Flux units are $\times 10^{-14} \text{ ergs cm}^{-2} \text{ s}^{-1}$.

upper limit we used 0.2 keV to determine a 0.2–10 keV flux of $4 \times 10^{-15} \text{ ergs cm}^{-2} \text{ s}^{-1}$ and an unabsorbed flux of $6 \times 10^{-15} \text{ ergs cm}^{-2} \text{ s}^{-1}$. The corresponding X-ray luminosity is $7 \times 10^{28} \text{ ergs s}^{-1}$ for an assumed distance of




FIG. 4.— *XMM-Newton*/EPIC pn and MOS data for both February and March data sets with best-fit thermal MEKAL+blackbody model. The discrepancies around 0.8 keV simply reflect the combination of the low spectral resolution and the complex of emission lines in the emitting plasma in this region.

300 pc.

4. CONCLUSIONS

Our X-ray observations of the extremely low accretion rate Polar SDSSJ1553 reveal a low count rate of $0.030 \pm 0.003 \text{ c/s}$, no detectable variation of the X-ray flux throughout the orbit (while the optical shows large variations due to the changing amplitude of the cyclotron harmonics), and a spectrum that can be modeled by a thermal plasma with a temperature near 0.8 keV. The X-ray flux translates into a luminosity below $10^{29} \text{ ergs s}^{-1}$. The low count rate of SDSSJ1324 (0.0012 ± 0.0003) is also consistent with this X-ray luminosity. All these characteristics imply an origin of the X-rays from coronal emission on the secondary star, rather than an accretion shock. The sensitivity of *XMM-Newton* has allowed the base level of X-ray emission from a close binary to be measured. Further X-ray monitoring of these stars whose mass transfer has turned off can reveal if the flare activity of M stars in close binaries is enhanced due to the close proximity of its companion white dwarf.

We acknowledge an anonymous referee for helpful suggestions on the evolutionary state. This work was funded by NASA XMM grants NAG5-12938 and NNG04GG66G and NSF grant AST-02-05875 to the University of Washington. This work is based on observations obtained with *XMM-Newton*, an ESA science mission with instruments and contributions directly funded by ESA member states and the USA (NASA).

Funding for the creation and distribution of the SDSS Archive has been provided by the Alfred P. Sloan Foundation, the Participating Institutions, the National Aeronautics and Space Administration, the National Science Foundation, the U.S. Department of Energy, the Japanese Monbukagakusho, and the Max Planck Society. The SDSS Web site is <http://www.sdss.org/>. The SDSS is managed by the Astrophysical Research Consortium (ARC) for the Participating Institutions. The Participat-

ing Institutions are The University of Chicago, Fermilab, the Institute for Advanced Study, the Japan Participation Group, The Johns Hopkins University, Los Alamos National Laboratory, the Max-Planck-Institute for Astronomy (MPIA), the Max-Planck-Institute for Astrophysics (MPA), New Mexico State University, University of Pittsburgh, Princeton University, the United States Naval Observatory, and the University of Washington.

REFERENCES

- Abazajian, K. et al. 2003, *AJ*, 126, 2081
 Abazajian, K. et al. 2004, *AJ*, in press
 de Martino, D. et al. 1998, *A&A*, 333, 31
 Fabbiano, G. 1982, *ApJ*, 262, 709
 Ferrario, L., Schmidt, G., Wickramasinghe, D. T. & Smith, P. 2002, poster at IAU Coll. 191, S. Africa
 Gänsicke, B. T., Beuermann, K., de Martino, D. & Thomas, H. C. 2000, *A&A*, 354, 605
 Kuijpers, J. & Pringle, J. E. 1982, *A&A*, 114, L4
 Malkov, O. Y., Piskunov, A. E. & Shpil’kina, D. A. 1997, *A&A*, 320, 79
 Mason, K. O. et al. 2001, *A&A*, 365, L36
 Mukai, K., Kinkhabwala, A., Peterson, J. R., Kahn, S. M., Paerels, F. 2003, *ApJ*, 586, 77
 Ramsay, G., Cropper, M., Wu, K., Mason, K. O. Cordova, F. A. & Priedhorsky, W. 2004, *MNRAS*, 350, 1373
 Szkody, P. et al. 2002, *AJ*, 123, 430
 Szkody, P. et al. 2003a, *AJ*, 126, 1499
 Szkody, P. et al. 2003b, *ApJ*, 583, 902
 Townsley, D. M. & Bildsten, L. 2004, *ApJ*, 600, 390
 Turner, M. et al. 2001, *MNRAS*, *A&A*, 365, L27
 Warner, B. in *Cataclysmic Variable Stars*, (Cambridge:CUP)
 Wickramasinghe, D. & Ferrario, L. 2000, *PASP*, 112, 873
 Woelk, U., & Beuermann, K. 1992, *A&A*, 256, 498
 Woelk, U., & Beuermann, K. 1993, *A&A*, 280, 173
 York, D. G. et al. 2000, *AJ*, 120, 1579